



2x qPCR Probe Master Mix - high ROX

Article	Content
SL-9802HR-smp	1 ml, 100 rxn × 20 µl
SL-9802HR-5ML	5 x 1 ml, 500 rxn × 20 µl
SL-9802HR-10ML	10 x 1 ml, 1000 rxn × 20 µl
SL-9802HR-20ML	20 x 1 ml, 2000 rxn × 20 µl



Long-Term Storage
at -20°C in the dark

Short-Term Storage
at 4°C in the dark

DESCRIPTION

Our **primaQUANT PROBE 2x qPCR Master Mix** is an optimized ready-to-use mixture for probe-based assays such as Taqman®, Beacons and MGBs. It contains a modified HotStart DNA Polymerase, as well as dNTPS and MgCl₂. The sophisticated buffer system provides fast kinetics and target amplification even for difficult templates.

The **primaQUANT 2x Master Mix** contains all components - you just need to add primers and template DNA/cDNA.



DID YOU KNOW?

- **primaQUANT PROBE** is also offered as a blue-colored mix for better handling.
- This Master Mix contains **high ROX** at a final concentration of 500 nM.
- For qPCR cyclers requiring other concentrations of ROX, **primaQUANT PROBE** is also available with low concentrations or without ROX.



Recommended Reaction Mixture per Well



BEFORE YOU START

- After thawing, please **invert the Master Mix tube 6-8 times**.
- **Do not vortex** the Master Mix to avoid damaging the enzyme.

Component	Stock Concentration	20 μ l Reaction	10 μ l Reaction	Final Concentration
2x primaQUANT Master Mix	2x	10 μ l	5 μ l	1x
Forward Primer	4 μ M	1 μ l	0.5 μ l	200 nM (100 - 400 nM recommended)
Reverse Primer	4 μ M	1 μ l	0.5 μ l	200 nM (100 - 400 nM recommended)
Probe	8 μ M	1 μ l	0.5 μ l	400 nM (200 - 600 nM recommended)
Template DNA	-	variable	variable	0.1 - 10 ng per reaction
Sterile Water	-	adjust to 20 μ l	adjust to 10 μ l	-



NOTE

For maximum efficiency and specificity, adjustments of annealing temperature as well as extension time, primer/probe concentration and template concentration may be needed.

CALCULATOR TOOL

Please feel free to download our Excel sheet calculator to calculate the necessary volumes:
calculator.steinbrenner-laborsysteme.de



qPCR
KnowledgeCenter

Standard Protocol



NOTE

- For the majority of qPCR assays, standard cycling conditions can be applied for the majority of qPCR assays out-of-the box.
- However, cycling conditions strongly depend on the primer, probe, amplicon and input material and thus some of these factors might need adjustments.

3-STEP PROTOCOL

Step	Time	Temperature	
Initial Denaturation	1 - 3 minutes	92°C - 95°C	
Denaturation	5 - 10 seconds	92°C - 95°C	
Annealing	1 - 5 seconds	60°C depending on primer	25 - 40 cycles
Extension	10 - 20 seconds	72°C	

2-STEP PROTOCOL

Step	Time	Temperature	
Initial Denaturation	1 - 3 minutes	92°C - 95°C	
Denaturation	5 - 10 seconds	92°C - 95°C	
Annealing / Extension combined	10 - 20 seconds	60°C depending on primer	25 - 40 cycles

Ultra-fast Protocol



NOTE

- Ultra-fast cycling conditions highly depend on the ramping rate of your qPCR cycler, primer, probe, amplicon and input material and thus might need adjustments.
- Ultra-fast cycling conditions can be applied for the majority of qPCR assays out-of-the box, provided that your primer/probe sets do not show unspecific binding.

3-STEP PROTOCOL

Step	Time	Temperature	
Initial Denaturation	1 minute	92°C - 95°C	
Denaturation	1 - 5 seconds	92°C - 95°C	25 - 40 cycles
Annealing	1 - 5 seconds	60°C depending on primer	
Extension	1 second	72°C	

2-STEP PROTOCOL

Step	Time	Temperature	
Initial Denaturation	1 minute	92°C - 95°C	
Denaturation	1 second	92°C - 95°C	25 - 40 cycles
Annealing / Extension combined	1 - 5 seconds	60°C depending on primer	



Applications

Probe-based quantitative PCR

- TaqMan® Probes
- Any kind of Dual-Labeled Hydrolysis Probe
- Molecular Beacons
- Scorpion Probes

DNA Genotyping

DNA SNP Analysis

RNA and miRNA Expression

Multiplexing (up to 4 colors)

Transcript Variant Analysis

QUALITY CONTROL PROCEDURE

Our **primaQUANT PROBE 2x qPCR Master Mix** undergoes stringent quality controls. Each lot is tested in a probe-based qPCR with cDNA and lambda DNA input.

Enzyme purity and homogeneity of > 98 % is validated using a Bioanalyzer SDS protein electrophoresis.

All **primaQuant** Master Mixes are free of detectable endonuclease- & exonuclease activity:

- Incubation of 1 µg of plasmid DNA with 5 U for 4h at 37°C and 72°C
- Incubation of 1 µg of a DNA size standard with 5 U for 4h at 37°C and 72°C

For more information, please visit our website: www.steinbrenner.de



Steinbrenner
Laborsysteme GmbH

In der Au 17 | 69257 Wiesenbach

+49 (0) 6223 / 96 73 00

mail@steinbrenner.de



Further Products

Products that may also interest you

REVERSE TRANSCRIPTION KIT



For the efficient cDNA synthesis out of total RNA extractions try the primaREVERSE RT-Kit with article number SL-9540.

